



Study of Physicochemical parameters in Water

Sandeep Kumar

Research Scholar, Department of Chemistry JJT UNIVERSITY

Abstract : Water testing is essential before drinking, washing, or using in agriculture or industry. Multiple physicochemical factors should be considered while evaluating water. The purpose for which we will be using the water, as well as the desired level of quality and purity, will determine the criteria we use to test it. Impurities in water may be in several states: suspended, dissolved, microbiological, or even bacterial. Chemical tests, including those for Physical tests like temperature, color, odor, pH, turbidity, TDS, and so on must be combined with chemical tests like BOD, COD, dissolved oxygen, alkalinity, hardness, and so on.

The only way to ensure that you're getting the purest water possible is to have it tested for naturally occurring contaminants like pesticides, heavy metals, and even traces of steel. Drinking water should naturally be safe and have the appropriate minerals. For the most part, these statistics are tracked carefully in advanced nations.

Key Words : Water, temperature, color, odor, pH, turbidity, TDS, and so on must be combined with chemical tests like BOD, COD, dissolved oxygen, alkalinity, hardness etc.

Introduction

Commonly referred to as water, this material has a molecular formula of H₂O and consists of two hydrogen and one oxygen atoms. All other forms of life depend on it. In spite of the fact that its liquid form is the one most often thought of, water may also exist in solid (ice) and gaseous (vapour) forms (steam or water vapour). In addition to solid water, liquid-crystal water may be found on surfaces with a high affinity for water.

1.460 1021 kilogrammes (pt) of water (71% of Earth's surface) is found in the world's seas and other big bodies of water. 1.6% of subsurface aquifers and 0.001% of cloud water are also affected (these Formation occurs from suspended solid and liquid particles of water in the air). Saltwater oceans account for 97% of the world's total, with the remaining 3% coming from freshwater rivers, lakes, and ice caps. The ponds also contain water, accounting for 0.6% of the total. Only a fraction of the water on Earth is stored in containers, organisms, goods, and food. Ice-capped mountains, glaciers, underground water sources, and frozen lakes sometimes provide Earth with drinkable quantities of water.



Temperature: In a stable system, the water's temperature determines the rate of all chemical processes and has an effect on fish reproduction, growth, and resistance to disease. Fish are very sensitive to temperature fluctuations and might perish if the water suddenly warms or cools dramatically.

PH: The PH scale is the most reliable indicator of water's corrosiveness. The more corrosive water is, the lower its pH value. Both electric conductivity and overall alkalinity increased as the pH value increased. Low oxygen levels correspond with increased temperatures during the summer months, which may be responsible for a decrease in photosynthetic activity absorption of carbon dioxide and bicarbonate. The pH of water may change as a result of exposure to a wide range of substances. The shift in physicochemical kingdom is having a greater effect on the stability of carbon dioxide, carbonate, and bicarbonate, as shown by the increased pH values found.

(Electrical conductivity): There is a strong relationship between conductivity and nine other measures of water quality. A number of factors are considered, including temperature, pH, alkalinity, total hardness, calcium, overall solids, total dissolved solids, chemical oxygen demand, chloride awareness, and iron awareness. Researchers Navneet Kumar et al. (2010) proposed a technique for determining the quality of underground drinking water in the region of investigation by lowering the conductivity of the water; this approach may be applicable to the management of water quality in other places. An EC meter may be used to measure the water resistance between two plated electrodes. The conductance values from a standard KCl solution were used to evaluate the precision of the measuring device.

Carbon dioxide: All aquatic settings produce carbon dioxide as a byproduct of decomposing organic matter, and the degree to which this gas varies depends largely on the metabolic activity of the internet's atmosphere. Therefore, while conducting research on the biogeochemistry of aquatic environments, it is necessary to prioritize the characteristics of the carbon dioxide apparatus. One good illustration of this would be the fact that carbon dioxide (CO₂) is the most important greenhouse gas on Earth. It is a method that may be used to measure the environmental output or metabolism of an aquatic system as it passes via the air-water or sediment-water interface, and it is one of the most serious topics that can be found in the study of international commerce. An aqueous carbon dioxide device makes it very simple to acquire readings of a variety of parameters, including pH (pCO₂), measurements of total alkalinity and dissolved inorganic carbon (TA). A photometer may be used to measure the concentration of



carbon dioxide (pCO_2) in the water's surface., and the amount of dissolved inorganic carbon CO_2 can be determined with the assistance of a kilometer or infrared CO_2 analyzer. The measurement of alkalinity is accomplished by titrating a sample of CO_2 water with HCl until the point when the CO_2 equivalence point is reached.

Alkalinity: Alkalinity serves as a pH buffer and is mostly made up CO_3^{2-} and HCO_3^- (HCO_3^-). Alkalinity, pH, and hardness of water all have a role in determining how dangerous certain compounds in water really are. Using phenolphthalein and methyl orange as indicators, an HCl titration quickly and easily provides the answer. Hydroxyl and carbonate ions are the primary causes of alkalinity in boiler water. Boiler water has to have a high hydroxyl alkalinity to prevent corrosion. Problems with operation, such as foaming, may be caused by excessive corrosiveness. Attenuation, a kind of caustic assault of the boiler caused by very high caustic stages, is possible.

Dissolved Oxygen: One of the most crucial factors is dissolved oxygen because of the information it provides about things like bacterial activity, photosynthesis, nutrient availability, stratification, and more. A rise in temperature and an increase in microbial activity may lead to a drop in dissolved oxygen levels later in the summer. Rising temperatures and periods of intense sunlight increase the percentage of dissolved gases (O_2 and CO_2), leading to higher levels of dissolved oxygen in the summer. Summer's long days and strong sunlight cause phytoplankton to produce more oxygen and use more carbon dioxide. This is likely the cause of the summertime increase in O_2 levels. After 5 days in 293 K, Titrimetric analysis utilizing Winkler's technique is used to determine the amount of dissolved oxygen in the sample.

Chemical Oxygen Demand : Some other measure of organic count pollution in water is chemical oxygen demand, and it is expressed in milligrammes per litre. The chemical oxygen demand is the amount of dissolved oxygen required for the chemical oxidation of organic molecules in water, and like the biological oxygen demand (BoI), it is an essential indicator of the environmental health of ground water components. They are often used for treating wastewater but are seldom used to water in general.

Sulfate: The nephelometric method is used to determine the level of turbidity by comparing the measured level of visibility to the known level of visibility of an artificially generated sulphate solution. Because barium sulphate is used to create turbidity, Turbidity may be avoided using barium chloride, and it can be created with a mixture of organic materials (glycerol or gum acetia) and sodium chloride.



Ammonia: (Nitrogen) A spectroscopic colorimetric measurement is made by mixing Nestler's reagent with 425 nm radiation to form a colour complex. Alkaline reaction conditions significantly alter the hardness of the water.

For pH values greater than 12, calcium is determined via complexometric titration utilising a modern EDTA solution and Patton and Reader's indicator. Sodium hydroxide, 4N concentration, is added to achieve these conditions. Calcium in a sample can be detected by plotting the titer (EDTA solution concentration) against a known sample volume.

Biochemical oxygen call for: The amount of natural bacterial infection in water is expressed as a biochemical oxygen demand in millimoles per litre. Oxygen demand in biochemistry refers to the quantity of oxygen in solution essential in biological reactions involving the breakdown of organic molecules and the oxidation of certain inorganic components (eg, iron, sulfites).essential in biological reactions involving the breakdown of organic molecules and the oxidation of certain inorganic components (eg, iron, sulfites). Standard biochemical oxygen demand testing takes place over the course of five days.

Magnesium: Black tea brewed with eriochrome as an indicator is used in a complicated metric titration with a common solution of EDTA at a buffer pH of 10.0. Ammonium hydroxide and ammonium chloride combine to form the buffer solution. It is difficult to change the solution's pH by titrating it.

Sodium: Help from a flame photometer is used to get an accurate reading. The instrument is calibrated using salt solutions of known concentrations (from 1 to 100 mg/l). The dilution effect is applied to the data after very concentrated samples have been adequately diluted with distilled water.

Silicates and Phosphates: These are also measured spectroscopically. Under highly acidic conditions, the molybdate ion becomes yellow as phosphate and silicate ions move over it. There is a direct correlation between the concentrations of phosphate and silicate in the composition and the resulting hue. Ascorbic acid and tartaric acid (potassium antimonyl tartrate) are examples of weak reducing marketers that can be used to deactivate phosphate complexes, while hydrazine and bisulphite are required for silica complexes. The simplified structure has a light blue hue.

Chloride: To do so, you'll need some Use an indicator solution, Titrate a known volume of your sample against a standard silver nitrate solution using a reagent such as potassium chromate in water or eosin/fluorescein in alcohol.Adsorption indicators are different from



precipitation indicators like the former bureaucracy's crimson silver substance, which is precipitated out of a chloride solution.

Total Hardness:

Total hardness growth boiling point of water its far soap eating ability of water sample. Calcium and magnesium cations, or cat ions, are the primary precipitates responsible for hardness. A variety of other cat ions, including strontium, iron, and manganese, also contribute to the hardness of water. The important anions which can be answerable for hardness are carbonates, bicarbonates, sulphate chlorides silicate and nitrate.

Due to the abundance of rocks and minerals, the number of divalent cations in water is the single most important factor in establishing its overall hardness. Ca^{++} and Mg^{++} and the polyvalent cations Al^{3+} , Ba^{2+} , Fe^{2+} , Mn^{2+} , Sr^{2+} and Zn^{2+} . A large quantity of soap and detergent may be used in hard water since it does not generate suds when combined with the soap. The titrimetric technique is used to get the value in milligrammes of CaCO_3 per millilitre. There are weak associations between cardiovascular fitness and water hardness, according to the literature". Improvements in children's eczema have been linked to the use of hard water systems at home. Hardness descriptions are roughly proportional to mineral content:

Water Classification Based on Hardness

Water Class	Hardness as CaCO_3(mg/L)
Soft	0–75 mg/L
Moderately hard	75 –150 mg/L
Hard	150 –300 mg/L
Very hard	>300 mg/L

High levels of hardness were found in the research region, and the obvious cause is the inappropriate or nonexistent disposal of sewage and industrial pollutants. If sulphate ions are present, drinking hard water can cause diarrhea.

Dissolved Oxygen (DO):

The amount of dissolved oxygen is a critical factor in identifying top-tier water. Evidence that physical and biological processes dominate in aquatic environments. Oxygenated waters taste the most delicious. The same old DO in water is 7.6 mg/L at 30°C and 7.0 mg/L at 35°C. Oxygen in water comes from the photosynthetic activities of aquatic plants, phytoplankton,



and aeration, however the amount is decreasing due to many organic contaminants. Its volume is determined by use of an iodometric technique known as the Winkler Azide Modification, and an instrumentation and analyzer package are used to precisely quantify the result. If the concentration of oxygen in water drops, normal populations will be disrupted, and anaerobic microorganisms will proliferate.

One crucial factor in water testing is the amount of oxygen that has been dissolved in the liquid. A combination of chemical, organic, and biological activities in the water are responsible for establishing the dissolved oxygen concentration. Atmospheric aeration and photosynthesis contribute to the oxygen content of water, while oxygen is used up in respiration, the breakdown of organic matter, and evaporation. In aquatic environments, the level of dissolved oxygen in the water is a good indicator of the health of the ecosystem, and its concentration may be quantified as a function of both the time and depth at which an inspection is conducted.

Biochemical Oxygen Demand (BOD):

Oxygen, the Vital Nutrient The quantity of Oxygen that microorganisms use in maintaining the organic stability is the quantity that is needed. In aquatic equipment, the amount of oxidizable organic matter is what determines the BOD cost. DO and BOD have opposite temporal relationships. Sixty-four more inorganic compounds mean more money out of pocket for BOD treatment. The thin-cubating method is used to get this value at 20 degrees Celsius.

Chemical Oxygen Demand (COD):

Chemical oxygen Call for is quantity of oxygen required by organic depend in a pattern of water for its oxidation by robust chemical oxidizing marketers like potassium dichromate it's miles expressed in pm or mg/L. COD fee measurement is vital as wherein BOD check can not carried out appropriately due to presence of toxin or different damaging condition for growth of bacteria. The downside of this test is that some natural fabric like benzene, pyridine etc are not oxidized by means of dichromate techniques and some inorganic fabric like nitriles, sulphides, thiosulphate, reduced metal ions eat oxygen for their oxidation. The study at can't provide concept about whether waste is degradable biologically and price in their oxidation. In preferred COD check is beneficial than BOD test for business waste.

Sulphate:

These contaminants have become synonymous with polluted water. Sources of SO₄²⁻ in surface and subsurface water include industrial effluents, household wastes, chemical fertilizers, as well as the decomposition of minerals including barite (BaSO₄), epsomite



($\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$), and gypsum ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$). The results are measured in milligrammes per litre (mg/L) to account for turbidity. Increased sulphate concentrations in water have been linked to digestive system problems, a laxative or cathartic effect on humans, and an increased risk of renal and heart disease.

Phosphate:

Orthophosphate, natural phosphate, and polyphosphate are all forms of phosphorus found in nature. In addition to being released into the environment with the help of fertilisers and detergents, phosphates also act as a nutrient pollutant in water, contributing to issues like eutrophication, algal blooms, and significant diurnal variations in dissolved oxygen. It's the main component that makes up things like calcium and phosphorus in bones. High phosphorus levels may potentially contribute to bone disorders.

Nitrate:

NO_3 and NO_2 are two of the most common nitrogen anions, and they may be found in both underground and aboveground water supplies. An rise in the quantity of nitrogen-containing compounds in the water supply is a consequence of both the extensive use of nitrogenous fertilizer on farms and the disposal of human and animal waste into ground water. Seepage also contaminates water supplies. Extreme levels of NO_3 and NO_2 . The lack of oxygen in the blood may lead to methemoglobinemia and blue baby syndrome in infants, and it can induce stomach cancer in adults.

Chloride:

The anion component of table salt, chloride, is the predominant particle in the human diet. Without being altered in any way, it is expelled from the body via the digestive system. Chloride concentrations in surface and groundwater are caused by a variety of human and natural activities, Water pollution is caused by a wide variety of human activities and natural processes, including road salt runoff, inorganic fertiliser use, landfill leachates, septic tank effluents, industrial effluents, sea water intrusion in coastal areas, and irrigation drainage. Chloride ingestion at concentrations as high as 2,000 mg/L does not present any dangers to human health and has only a small impact on the way things taste. When chloride is added to water, the corrosiveness of the water as well as its electrical conductivity increases, which both result in increased concentrations of steel in the water.

Fluoride:



Having enough fluoride in your diet is important for both animals and humans. Fluoride occurs naturally in a number of minerals, including fluorite, topaz, fluorapatite, crylite, mica, and others, and enters water as a result of weathering. Bone metabolism is aided by this, and dental fluorosis, skeletal fluorosis, and fluorosis of the teeth are all avoided thanks to the reduction of fluoride levels in the water we drink. This may be measured in milligrammes per millilitre using a specific ion electrode.

Sodium:

Sodium controls blood pressure and volume, helps keep fluids in balance, transmits nerve impulses, and affects muscle contraction and relaxation, making it an essential ingredient for all animal life. Sodium salts are very soluble in water and their concentrations increase due to both natural and human-made processes. There are numerous different signs of salt overload, but two of the most serious are heart failure and high blood pressure. Nausea, vomiting, seizures, muscle twitching, and tension are all possible short-term adverse effects.

Potassium:

Potassium is an alkaline metal that may be found naturally. It has a great resilience to weathering, and less of an impact occurs in water than Na. People with kidney diseases or other disorders, which includes heart disease, coronary artery disease, high blood pressure, diabetes, adrenal insufficiency, and preexisting hyperkalaemia, may wish to experience serious fitness repercussions from elevated potassium.

Calcium:

Calcium is found in large amount in rocks water get contaminated with calcium with the aid of reaching of calcium from rock. Calcium is maximum common metal found in natural water. The quantity of calcium in water can also range from 10 to 100mg/L the amount of calcium relies upon upon the nature of rock from which it's far contaminated another critical source of calcium commercial waste or sewage water. Calcium is chargeable for hardness of water. Calcium is crucial nutrient required via organism. It has no unsafe impact on human health. But higher amount of calcium in water is dangerous in industrial and residence maintain makes use of. High attention of calcium in water isn't makes use of.

Magnesium:

Magnesium, like calcium, is an essential element that may be found in most natural water sources. Magnesium is found in water from several types of rocks, as well as from sewage and industrial waste, although at a lower concentration than calcium. In other words, it makes water



harder. Natural water softening occurs when water percolates through soil and exchanges sodium ions. Magnesium does not have any known harmful effects in humans. Intense focus may be cleansing and dehydrating at first, but regular users quickly develop a resistance to its negative effects.

Metallic Elements:

Heavy metals are crucial to the functioning of all living things. We need Fe, Co, Cu, Mn, Mo, and Zn to function properly. Extreme values might be harmful to the living thing. Mercury, cadmium, lead, and chromium, to name a few, are toxic to humans and the environment; zinc and tin may corrode metals; and there are others that are harmful in various ways (e.G. As can also pollute catalysts). Miniscule amounts of several of these are required for human survival (Co, Cu, Cr, Mn, Ni). The disturbingly high levels of heavy metals in the environment are the result of both commercial interest and natural releases. The toxicity of trace metals is a major issue because of their pervasiveness and broad usage in contemporary life.

Iron:

Iron may be found naturally in the environment due to geological formations, acid drainage, and effluent discharges. Every living thing on Earth has traces of iron, making it an essential element. The considerable potential for organic toxicity of iron ions contributes to the stringent regulation of iron distribution in mammals. Iron excess leads to hemochromatosis and haemosiderosis.

Manganese:

Surface and groundwater resources contain manganese naturally, especially in anaerobic or low oxidation conditions. The syndrome referred to as manganism related to impaired motor competencies and cognitive problems, excessive results at the kidneys and liver.

Zinc:

When it comes to plants, animals, and bacteria, zinc is an essential trace element. Wastes and natural phenomena provide clear evidence of this. As a result of Zn's presence in water, it becomes bitter and astringent. Zinc fumes are very toxic, and being exposed to them may cause a wide range of unpleasant symptoms in humans, such as respiratory distress, fever, chills, gastrointestinal distress, abdominal pain, vomiting, lethargy, depression, and even paralysis of the lower and upper regions of the brain. Retardation, immaturity, and anaemia are exacerbated by a lack of zinc, as are recurrent illnesses, lowered immunity, hypozincemia, diarrhoea, pneumonia, acne, loss of taste, smell, memory, and hair, and skin lesions.



References :

1. DHIRENDRA MOHAN JOSHI ETAL (2009) studies on physicochemical parameters to assess the water quality of river ganga for drinking purpose in Haridwar district, Volume No.2, Issue No.1, PP no.105-203.
2. D. P. GUPTA ETAL (2009) Physiochemical Analysis of Ground Water of Selected Area of Kaithal City (Haryana) India, Volume No.1, Issue no.2, PP no.1-5.
3. KAIZER, A.N ETAL (2010) Physicochemical Characteristics and Heavy Metal Levels in Water Samples from Five River Systems in Delta State, Nigeria, Volume No.14, Issue No.1, PP No.83-87, ISSN 1119-8362.
4. GABRIEL I. OBIEFUNA ETAL (2010) Physicochemical Characteristics of groundwater quality from Yola Area, Northeastern Nigeria, Volume No.14, Issue No.1, PP No.5-11, ISSN 1119-8362.
5. ANURAG TEWARI ETAL (2010) A study on physical and chemical make-up of ground water quality, Volume No.2, Issue No.2, PP No.510-518, ISSN No: 0975-7384.
6. S. PRADHAN ETAL (2011) Characterization of Groundwater Quality for Irrigation in 'Gohana' Block of Sonapat District, Haryana, Volume No.1, Issue No.2, PP No.63-70, ISSN 0973-032X.
7. DATTATRAYA BHART ETAL (2011) Physical and chemical make-up of Bore well water quality in Nagpur region (South zone), Volume No.3, Issue No.2, PP no.922-927, ISSN No: 0975-7384.
8. SULEKH CHANDRA ETAL (2011) Evaluation of Physicochemical Characteristics of Various River Water in India, Volume No.8, Issue No.4, PP No.1-11, ISSN: 0973-4945.
9. EZERIBE, A. I ETAL (2012) chemical and physical properties of well water samples from some villages in Nigeria with cases of stained and mottle teeth, Volume No.7, Issue No.1, PP No.1-3, ISSN 1597-6343.
10. SULTAN SINGH ETAL (2012) Groundwater Quality Analysis of Safidon and Julana Blocks of District Jind, Haryana, India, Volume No.4, Issue No.2, PP No.1-10.